arXiv:0902.4869v1 [math.FA] 27 Feb 2009

QUANTUM ERROR CORRECTION AND HIGHER RANK NUMERICAL RANGES OF NORMAL MATRICES

HWA-LONG GAU, CHI-KWONG LI, YIU-TUNG POON, AND NUNG-SING SZE*

Abstract. The higher rank numerical range is useful for constructing quantum error correction code for a noisy quantum channel. It is known that if a normal matrix $A \in M_n$ has eigenvalues $a_1, \ldots, a_n$, then its rank-k numerical range $\Lambda_k(A)$ is the intersection of convex polygons with vertices $a_{j_1}, \ldots, a_{j_{n-k+1}}$, where $1 \le j_1 < \cdots < j_{n-k+1} \le n$. In this paper, it is shown that the higher rank numerical range of a normal matrix with m distinct vertices can be written as the intersection of no more than m closed half planes. In addition, given a convex polygon $\mathcal{P}$ a construction is given for a normal matrix $A \in M_n$ with minimum n such that $\Lambda_k(A) = \mathcal{P}$. In particular, if $\mathcal{P}$ has p vertices, with $p \ge 3$, there is a normal matrix $A \in M_n$ with

$$n \le \max\{p + k - 1, 2k + 2\}$$

such that $\Lambda_k(A) = \mathcal{P}$.

1. Introduction

Let M_n be the algebra of $n \times n$ complex matrices regarded as linear operators acting on the n-dimensional Hilbert space $\mathbb{C}^n$. In the context of quantum information theory, if the quantum states are represented as matrices in M_n, then a *quantum channel* is a trace preserving completely positive map $L : M_n \to M_n$ with the following operator sum representation

$$L(A) = \sum_{j=1}^{r} E_j^* A E_j,$$

where $E_1, \ldots, E_r \in M_n$ satisfy $\sum_{j=1}^{r} E_j E_j^* = I_n$. The matrices $E_1, \ldots, E_r$ are known as the *error operators* of the quantum channel L. A subspace V of $\mathbb{C}^n$ is a *quantum error correction code* for the channel L if and only if the orthogonal projection $P \in M_n$ with range space V satisfies $PE_i^*E_jP = \gamma_{ij}P$ for all $i, j \in \{1, \ldots, r\}$; for example, see [7, 8, 9]. In this connection, for

---

2000 *Mathematics Subject Classification.* 15A60, 15A90, 47N50, 81P68.

Key words and phrases. Quantum error correction, higher rank numerical range, normal matrices, convex polygon.

Research of Gau was supported by National Science Council of the Republic of China. Research of Li was supported by USA NSF and the William and Mary Plumeri Award.

*Corresponding author.

$1 \leq k < n$ researchers define the *rank-k numerical range* of $A \in M_n$ by

$$\Lambda_k(A) = \{\lambda \in \mathbb{C} : PAP = \lambda P \text{ for some rank-}k \text{ orthogonal projection } P\},$$

and the *joint rank-k numerical range* of $A_1, \ldots, A_m \in M_n$ by $\Lambda_k(A_1, \ldots, A_m)$ to be the collection of complex vectors $(a_1, \ldots, a_m) \in \mathbb{C}^{1\times m}$ such that $PA_jP = a_jP$ for a rank-k orthogonal projection $P \in M_n$. Evidently, there is a quantum error correction code V of dimension k for the quantum channel L described above if and only if $\Lambda_k(A_1, \ldots, A_m)$ is non-empty for $(A_1, \ldots, A_m) = (E_1^*E_1, E_1^*E_2, \ldots, E_r^*E_r)$. Also, it is easy to see that if $(a_1, \ldots, a_m) \in \Lambda_k(A_1, \ldots, A_m)$ then $a_j \in \Lambda_k(A_j)$ for $j = 1, \ldots, m$.

When $k = 1$, $\Lambda_k(A)$ reduces to the *classical numerical range* defined and denoted by

$$W(A) = \{x^*Ax \in \mathbb{C} : x \in \mathbb{C}^n \text{ with } x^*x = 1\},$$

which is a useful concept in studying matrices and operators; see [6]. Recently, interesting results have been obtained for the rank-k numerical range and the joint rank-k numerical range; see [1, 2, 3, 4, 5, 11, 12, 13, 14, 16]. In particular, an explicit description of the rank-k numerical range of $A \in M_n$ is given in [14], namely,

$$(1.1)\quad \Lambda_k(A) = \bigcap_{\xi\in[0,2\pi)} \{\mu \in \mathbb{C} : e^{-i\xi}\mu + e^{i\xi}\overline{\mu} \leq \lambda_k(e^{-i\xi}A + e^{i\xi}A^*)\},$$

where $\lambda_k(X)$ is the kth largest eigenvalue of a Hermitian matrix X.

In the study of quantum error correction, there are channels such as the randomized unitary channels and Pauli channels whose error operators are commuting normal matrices. Thus, it is of interest to study the rank-k numerical ranges of normal matrices.

Given $S \subseteq \mathbb{C}$, let $\operatorname{conv} S$ denote the smallest convex subset of $\mathbb{C}$ containing S. For a normal matrix $A \in M_n$ with eigenvalues $a_1, \ldots, a_n$, it was conjectured in [3, 4] that

$$(1.2)\qquad \Lambda_k(A) = \bigcap_{1\leq j_1<\cdots<j_{n-k+1}\leq n} \operatorname{conv}\{a_{j_1}, \ldots, a_{j_{n-k+1}}\}$$

is again a convex polygon including its interior (if it is non-empty). This conjecture was confirmed in [14] using the description of $\Lambda_k(A)$ in (1.1). *In our discussion, a polygon would always mean a convex polygon with its interior.*

In this paper, we improve the description (1.2) of the rank-k numerical range of a normal matrix. In particular, in Section 2 we show that for a normal matrix A with m distinct eigenvalues, $\Lambda_k(A)$ can be written as the intersection of no more than m closed half planes in $\mathbb{C}$. Consequently, if $\Lambda_k(A) \neq \emptyset$, then it is a polygon with no more than m vertices. We then consider the "inverse" problem, namely, for a given polygon $\mathcal{P}$, construct a normal matrix $A \in M_n$ with $\Lambda_k(A) = \mathcal{P}$. In other words, we study the necessary condition for the existence of quantum channels whose error operators have prescribed rank-k numerical ranges. It is easy to check that $\Lambda_k(\tilde{A}) = \mathcal{P}$ if $\tilde{A} = A \otimes I_k$ with $W(A) = \mathcal{P}$. Our goal is to find a normal matrix $\hat{A}$ with smallest size so that $\Lambda_k(\hat{A}) = \mathcal{P}$. To achieve this, we give a necessary and sufficient condition for the existence of a normal matrix $A \in M_n$ so that $\Lambda_k(A) = \mathcal{P}$ in terms of k-regular sets in $\mathbb{C}$ (see the definition in Section 3). Furthermore, we show that the problem of finding a desired normal A is equivalent to a combinatorial problem of extending a given p element set of unimodular complex numbers to a k-regular set. We then give the solution of the problem in Section 4. As a consequence of our results, if $\mathcal{P}$ is a polygon with p vertices, then there is a normal matrix $A \in M_n$ with

$$n \leq \max\{p + k - 1, 2k + 2\}$$

such that $\Lambda_k(A) = \mathcal{P}$. Moreover, this upper bound is best possible in the sense that there exists $\mathcal{P}$ so that there is no matrix of smaller dimension with rank-k numerical range equal to $\mathcal{P}$.

2. Construction of higher rank numerical ranges

The purpose of this section is to study the following.

Problem 2.1. Find an efficient way to construct $\Lambda_k(A)$ for a normal matrix $A \in M_n$ and determine/estimate the number of vertices of $\Lambda_k(A)$.

Suppose the eigenvalues of A are collinear. Then by a translation, followed by a rotation, we may assume that A is Hermitian with eigenvalues $a_1 \geq \cdots \geq a_n$. Then we have $\Lambda_k(A) = [a_{n-k+1}, a_k]$. So we focus on those normal matrix A whose eigenvalues are not collinear.

Let us motivate our result with the following example.

Example 2.2. Let $a_1, \ldots, a_n$ be the eigenvalues of A, with $n \geq 3$. If $\operatorname{conv}\{a_1, \ldots, a_n\} = \mathcal{P}$ is an n-sided polygon, then every a_j is an extreme point of $\mathcal{P}$. We may assume that $a_1, \ldots, a_n$ are arranged in the counterclockwise direction on the boundary of $\mathcal{P}$. For convenience of notation, we would let $a_j = a_{j-n}$ for $j > n$. Now for every j, let L_j be the line passing through a_j and a_{j+k} and $\mathcal{H}_j$ be the closed half plane determined by L_j which does not contain a_ℓ for $j < \ell < j + k$. We have $\Lambda_k(A) = \bigcap_{j=1}^{n} \mathcal{H}_j$.

One can apply (1.2) or Corollary 2.8 to verify the above example. Note that each $\mathcal{H}_j$ in Example 2.2 contains exactly $n - k + 1$ eigenvalues of A. Example 2.9 below is a special case.

In general, given any two distinct eigenvalues a_r and a_s of A, let $L(r, s)$ be the (directed) line passing through a_r and a_s. The closed half plane

$$H(r, s) = \{z \in \mathbb{C} : \operatorname{Im}\left((\bar{a}_s - \bar{a}_r)(z - a_r)\right) \geq 0\}$$

is called the *left closed half plane* determined by $L(r,s)$. In our discussion, it is sometimes convenient to write

$$H(r,s) = \{z \in \mathbb{C} : \mathrm{Re}\,(e^{-i\xi} z) \le \mathrm{Re}\,(e^{-i\xi} a_r)\}$$

with $\xi = \arg(a_s - a_r) - \pi/2$.

We have the following.

Theorem 2.3. *Suppose $A \in M_n$ is normal with eigenvalues $a_1, \dots, a_n$ such that $\Lambda_k(A)$ is a non-degenerate polygon. Then*

$$\text{(2.1)} \quad \Lambda_k(A) = \cap\{H(r,s) : a_r \ne a_s,\ H(r,s) \text{ contains at least } n-k+1 \text{ eigenvalues of } A \text{ (counting multiplicities) }\}.$$

If A has m distinct eigenvalues, then at most m distinct $H(r,s)$ are needed in the intersection in (2.1).

Proof. Let $\mathcal{S}$ be the set of index pairs (r,s) such that $H(r,s)$ contains at least $n-k+1$ eigenvalues of A. For each $(r,s) \in \mathcal{S}$, since $H(r,s)$ is convex and contains at least $n-k+1$ eigenvalues of A, we have

$$\Lambda_k(A) = \bigcap_{1 \le j_1 < \cdots < j_{n-k+1} \le n} \mathrm{conv}\,\{a_{j_1}, \dots, a_{j_{n-k+1}}\} \subseteq H(r,s).$$

It follows that

$$\Lambda_k(A) \subseteq \bigcap_{(r,s)\in\mathcal{S}} H(r,s). \tag{2.2}$$

To prove the reverse inclusion in (2.2) and also the last assertion, suppose A has m distinct eigenvalues $a_1, \dots, a_m$, and $\Lambda_k(A)$ is a non-degenerate polygon. For each $1 \le r \le m$, let $\mathcal{S}_r = \{s : (r,s) \in \mathcal{S}\}$. Then

$$\Lambda_k(A) \subseteq \bigcap_r \left(\bigcap_{s \in \mathcal{S}_r} H(r,s) \right).$$

For each r such that $\mathcal{S}_r \ne \emptyset$, $\bigcap_{s\in\mathcal{S}_r} H(r,s)$ has nonempty interior because $\Lambda_k(A)$ is non-degenerate. Therefore, there exists $s_r \in \mathcal{S}_r$ such that $a_{s_r} \in H(r,s)$ for all $s \in \mathcal{S}_r$. We are going to show that

$$\bigcap\{H(r,s_r) : 1 \le r \le m,\ \mathcal{S}_r \ne \emptyset\} \subseteq \Lambda_k(A). \tag{2.3}$$

Then (2.2) will ensure that the equality holds in (2.3).

Suppose $\mu \notin \Lambda_k(A)$. We **claim** that $\mu \in \mathbb{C} \setminus H(t, s_t)$ for some $t \in \{1, \dots, m\}$. To prove our claim, note that by (1.2) there is a non-degenerate polygonal disk $\mathcal{W}$, which is the convex hull of $n-k+1$ eigenvalues of A such that $\mu \notin \mathcal{W}$. We may replace A by $A - \nu I$ for a suitable $\nu \in \mathbb{C}$ and assume that 0 is an interior point of $\mathcal{W}$. We may relabel the eigenvalues of A and assume that $\mathcal{W}$ has vertices $a_1, \dots, a_p$ arranged in the counterclockwise direction, where $p \ge 3$. For convenience of notation, we will let $a_0 = a_p$ and

$a_j = a_{j-p}$ if $j > p$, and extend the definition of $H(t, t+1)$ to all integers t accordingly. Then

$$\mathcal{W} = \bigcap_{1 \le t \le p} H(t, t+1).$$

Now, we can express the set $\mathbb{C} \setminus \mathcal{W}$ as the disjoint union of $p+1$ conical regions of the form

$$H(t-1, t) \setminus H(t, t+1) = \operatorname{conv}\{L_{t+1} \cup L_t\} \setminus L_{t+1}, \quad t = 1, \ldots, p,$$

where

$$L_t = \{a_t + r(a_t - a_{t-1}) : r > 0\}.$$

Since $\mu \notin \mathcal{W}$, we may relabel the vertices $a_1, \ldots, a_p$ and assume that

$$\mu \in H(1,2) \setminus H(2,3).$$

Now, consider the half space $H(2, s_2)$. Since $\Lambda_k(A)$ is non-degenerate, a_{s_2} lies in the interior of $H(2,3)$. It follows that

$$\mu \in H(1,2) \setminus H(2,3) \subseteq \mathbb{C} \setminus H(2, s_2).$$

Thus, our claim is proved, and the result follows. □

Note that Theorem 2.3 may not hold if the convex set $\Lambda_k(A)$ is a subset of a line segment. In fact, if $A \in M_n$ is Hermitian with eigenvalues $a_1 \ge \cdots \ge a_n$ and $n \ge 2k$, then $\Lambda_k(A) = [a_{n-k+1}, a_k]$, which cannot be written as the intersection of half spaces of the form $H(r, s)$. So, the conclusion of Theorem 2.3 does not hold. For normal matrices with non-collinear eigenvalues, the last assertion of Theorem 2.3 may fail to hold even though the first part of the theorem is valid as shown in the following examples.

Example 2.4. Let $A = \operatorname{diag}(a_1, \ldots, a_5) = \operatorname{diag}(0, 0, 1, 1, i)$. Then

$$\Lambda_2(A) = [0,1] = H(1,3) \cap H(3,1) \cap H(5,1) \cap H(3,5)$$

and $\Lambda_2(A)$ cannot be written as an intersection of less than 4 half spaces $H(r, s)$. Therefore, the last assertion of Theorem 2.3 does not hold.

Example 2.5. Let $A = \operatorname{diag}(a_1, a_2, a_3, a_4) = \operatorname{diag}(1, -1, i, -i)$. Then

$$\Lambda_2(A) = \{0\} = H(1,2) \cap H(2,1) \cap H(3,4) \cap H(4,3)$$

and $\Lambda_2(A)$ cannot be written as an intersection of less than 4 half spaces $H(r, s)$. Therefore, the last assertion of Theorem 2.3 does not hold.

Inspired by the above examples, we have the following.

Theorem 2.6. *Suppose $A \in M_n$ is normal with non-collinear eigenvalues $a_1, \ldots, a_n$ such that $\Lambda_k(A)$ is a subset of a line segment. Let p be the smallest integer such that*

$$\Lambda_k(A) = \bigcap_{j=1}^{p} H(r_j, s_j)$$

for some $1 \leq r_j,\ s_j \leq n$. *Then* $p \leq 4$. *Moreover, we have the following.*

(a) *If* $\Lambda_k(A)$ *is a non-degenerate line segment, then* $p = 4$.

(b) *If* $\Lambda_k(A)$ *is a singleton, then* $p \in \{3, 4\}$.

(c) *If* $\Lambda_k(A) = \emptyset$, *then* $p \in \{2, 3\}$.

Proof. Suppose $a_1, \ldots, a_n$ are the eigenvalues of A.

(a) Suppose $\Lambda_k(A)$ is a non-degenerate line segment. Then $n \geq 2k$, see [1]. We may assume that $\Lambda_k(A) \subseteq \mathbb{R}$. Let

$$R_1 = \{j : \mathrm{Im}\,(a_j) > 0\},\ R_2 = \{j : \mathrm{Im}\,(a_j) = 0\},\ \text{ and } R_3 = \{j : \mathrm{Im}\,(a_j) < 0\}.$$

Then each of R_1 and R_3 has at most $k-1$ elements and at least one of them is non-empty. By (1.2), $\Lambda_k(A) \subseteq \mathrm{conv}\,\{a_r : r \in R_2 \cup R_3\}$. Therefore, R_2 has at least 2 elements r, s such that a_r and a_s are distinct points in $\mathbb{C}$. Let

$$\begin{aligned} a = &\ \max\{L(r_i, r_j) \cap \mathbb{R} : 1 \leq i < j \leq 3, r_i \in R_i,\ r_j \in R_j,\ (r_i, r_j) \in \mathcal{S}\}, \\ b = &\ \min\{L(r_j, r_i) \cap \mathbb{R} : 1 \leq i < j \leq 3, r_i \in R_i,\ r_j \in R_j,\ (r_i, r_j) \in \mathcal{S}\}. \end{aligned}$$

Choose $r_{i_j} \in R_{i_j}$ for $j = 1, \ldots, 4$ such that

$$a = L\left(r_{i_1}, r_{i_2}\right) \cap \mathbb{R} \quad \text{ and } \quad b = L\left(r_{i_3}, r_{i_4}\right) \cap \mathbb{R}.$$

Then we have

$$\Lambda_k(A) = [a, b] = H(r, s) \cap H(s, r) \cap H\left(r_{i_1}, r_{i_2}\right) \cap H\left(r_{i_3}, r_{i_4}\right).$$

Clearly, a non-degenerate line segment cannot be written as an intersection of less than four half spaces $H(r, s)$.

(b) Suppose $\Lambda_k(A)$ is a singleton. We may replace A by $A - \nu I$ for some suitable ν and assume that

$$\{0\} = \Lambda_k(A) = \bigcap_{(r,s) \in \mathcal{S}} H(r, s), \tag{2.4}$$

where $\mathcal{S}$ is defined as in the proof of Theorem 2.2. Evidently, in the intersection (2.4), we only need half spaces of the form

$$H(r_j, s_j) = \{z \in \mathbb{C} : \mathrm{Re}\,(e^{-i\xi_j} z) \geq 0\} = \mathcal{H}_j$$

for $j = 1, \ldots, p$. Since the intersection is a singleton, we have $p \geq 3$. We may assume that $0 \leq \xi_1 < \xi_2 < \cdots < \xi_p < 2\pi$. If $p \geq 5$, then either $\xi_3 - \xi_1$ or $\xi_5 - \xi_3$ is less than π. If $\xi_3 - \xi_1 < \pi$, then we have $\mathcal{H}_1 \cap \mathcal{H}_2 \cap \mathcal{H}_3 = \mathcal{H}_1 \cap \mathcal{H}_3$ and $\mathcal{H}_2$ can be omitted. Similarly, $\mathcal{H}_4$ can be omitted if $\xi_5 - \xi_3 < \pi$. Hence, we have $p \leq 4$. From the proof, we can see that $p = 4$ if and only if

$$\{H(r, s) : (r, s) \in \mathcal{S} \ \text{ and } \ 0 \in L(r, s)\} = \{\mathcal{H}_j : 1 \leq j \leq 4\}$$

and $0 \leq \xi_1 < \xi_2 < \pi \leq \xi_3 = \xi_1 + \pi < \xi_4 = \xi_2 + \pi$.

(c) If $\Lambda_k(A) = \emptyset$, then by Helly's Theorem [10, Theorem 24.9], there are three half spaces $H(r_j, s_j)$ for $j = 1, 2, 3$, such that $\cap_{j=1}^{3} H(r_j, s_j) = \emptyset = \Lambda_k(A)$. If the boundary of two of these half spaces are parallel lines, then the two half spaces must be disjoint so that intersecting the two half spaces is the empty set. Clearly, we need to intersect at least two half spaces $H(r,s)$ to get $\Lambda_k(A)$. Therefore, $p \in \{2,3\}$. $\square$

By Theorems 2.3 and 2.6, we have the following.

Corollary 2.7. *Suppose* $A \in M_n$ *is normal with* m *distinct eigenvalues. If* $\Lambda_k(A) \neq \emptyset$, *then* $\Lambda_k(A)$ *is a convex polygon with at most* m *vertices.*

Corollary 2.8. *Suppose* $A \in M_n$ *is normal such that* $W(A)$ *is an* n*-sided polygon containing the origin as its interior point. Let* $v_1, \dots, v_n$ *be the vertices of* $W(A)$ *having arguments* $0 \le \xi_1 < \cdots < \xi_n < 2\pi$. *If* $k < n/2$, *then* $\Lambda_k(A)$ *is an* n*-sided convex polygon obtained by joining* v_j *and* v_{j+k}, *where* $v_{j+k} = v_{j+k-n}$ *if* $j+k > n$.

By Theorem 2.3, it is easy to see that the boundary of $\Lambda_k(A)$ are subsets of the union of line segments of the form $\operatorname{conv}\{a_r, a_s\}$ such that a_r and a_s satisfy the $H(r,s)$ condition. However, it is not easy to determine which part of the line segment actually belong to $\Lambda_k(A)$. Here are some examples.

Example 2.9. Let $A = \operatorname{diag}(a_1, \dots, a_n) = \operatorname{diag}(1, w, w^2, \dots, w^{n-1})$ with $w = e^{2\pi i/n}$. Then for $k \le n/2$, we have $\Lambda_k(A) = \cap_{j=1}^n H(j, j+k)$, where $a_{j+k} = a_{j+k-n}$ if $j+k > n$, and only a small part of $\operatorname{conv}\{w^{j-1}, w^{j-1+k}\}$ lies in $\Lambda_k(A)$.



$\Lambda_2(A)$ with $n = 9$



$\Lambda_3(A)$ with $n = 9$

Example 2.10. Suppose $B = \operatorname{diag}(1, i, -1, -i, 2, 2i, -2, -2i, 3, 3i, -3, -3i)$. One can see from the figures that the eigenvalues $1, i, -1, -i$ are interior points of $\Lambda_2(B)$ while these eigenvalues are the vertices of $\Lambda_3(B)$.



$\Lambda_2(B)$



$\Lambda_3(B)$

Remark 2.11. *Note that if the eigenvalues* $a_1, \ldots, a_n$ *of a normal matrix* A *is given, one can easily construct* $\Lambda_k(A)$ *by the following algorithm from the discussion in this section.*

Case 1. *If the eigenvalues lie on a line* $\mathcal{L}$, *then arrange the points with real parts in descending order if* $\mathcal{L}$ *is not perpendicular to the real axis; otherwise, arrange the points with imaginary parts in descending order. Relabel the subscripts as* $a_1, \ldots, a_n$ *according to this ordering. If* $n \geq 2k-1$, *then* $\Lambda_k(A)$ *is the line segment joining* a_k *and* a_{n-k+1}, *which may degenerate to a point if* $a_k = a_{n-k+1}$. *If* $n < 2k-1$, *then it is the singleton* $\{a_k\}$ *if* $a_k = a_{n-k+1}$; *otherwise, it is the empty set.*

Case 2. *Suppose the eigenvalues do not lie on a straight line. For each distinct eigenvalue* a_r, *identify the set* $\mathcal{S}_r$ *and the point* a_{s_r}. *Determine the intersection of the half spaces* $\mathcal{H}(r, s_r)$, *which is* $\Lambda_k(A)$.

3. MATRICES WITH PRESCRIBED HIGHER RANK NUMERICAL RANGES

We study the following problem in this section.

Problem 3.1. Let $k > 1$ be a positive integer, and let $\mathcal{P}$ be a p-sided polygon in $\mathbb{C}$. Construct a normal matrix A with smallest size (dimension) such that $\Lambda_k(A) = \mathcal{P}$.

If $\mathcal{P}$ degenerates to a line segment joining two points a_1 and a_2. Then the smallest n to get a normal matrix with $\Lambda_k(A) = \mathcal{P}$ is $n = 2k$, if a_1 and a_2 are distinct and $n = k$ if $a_1 = a_2$. So we focus on the case when the polygon $\mathcal{P}$ is non-degenerate.

A natural approach to Problem 3.1 is to reverse the construction of $\Lambda_k(A)$ in Example 2.2. Suppose we have a non-degenerate p-sided polygon $\mathcal{P}$, with vertices, $v_1, \ldots, v_p$. Without loss of generality, we may assume that 0 is in the interior of $\mathcal{P}$ and $v_j = r_j e^{i\theta_j}$, where $r_j > 0$ and $0 \leq \theta_1 < \theta_2 < \cdots < \theta_p < 2\pi$. Let L_j be the line passing through v_j and v_{j+1}. Suppose the following conditions hold.

1. $k < p/2$.
2. for every $1 \le j \le p$, L_j and L_{j+k} intersects at a_j.
3. $a_1, \dots, a_p$ are vertices of a p-sided convex polygon arranged in the counterclockwise direction around 0.

Let A be a normal matrix with eigenvalues $a_1, a_2, \dots, a_p$. Then $\Lambda_k(A) = \mathcal{P}$. The above construction fails when one or more of conditions (1) – (3) are not satisfied. Conditions (2) and (3) motivate the following definition.

Definition 3.2. Let $\Omega = \{z \in \mathbb{C} : |z| = 1\}$. Given distinct $\alpha_1,\ \alpha_2 \in \Omega$, $\alpha_2/\alpha_1 = e^{i\theta}$ for a unique $0 < \theta < 2\pi$. Then $[\alpha_1, \alpha_2] = \{e^{it}\alpha_1 : 0 \le t \le \theta\}$ is the arc on Ω from α_1 to α_2 in the counterclockwise direction. Let $(\alpha_1, \alpha_2) = [\alpha_1, \alpha_2]\backslash\{\alpha_1, \alpha_2\}$. The value θ is called the length of these intervals. Suppose $1 \le k \le n$ and $\Pi \subseteq \Omega$. Then Π is said to be k-regular if for each $\alpha \in \Pi$, $(\alpha, -\alpha) \cap \Pi$ contains at least k elements.

If $\Pi = \{e^{i\xi_j} : 1 \le j \le n\}$ with $0 \le \xi_1 < \cdots < \xi_n < 2\pi$, then the above definition is equivalent to

$$\sin(\xi_{r+k} - \xi_r) > 0 \quad \text{for all} \quad r = 1, \dots, n. \tag{3.1}$$

Here, for notational convenience, we set $\xi_m = \xi_{m-n}$ if $m > n$. For this reason, a set $\{\xi_1, \dots, \xi_n\} \subseteq [0, 2\pi)$ is also called k-regular if $\{e^{i\xi_j} : 1 \le j \le n\}$ is k-regular as defined in Definition 3.2. For $\xi, \xi' \in [0, 2\pi)$, $[\xi, \xi']$ will denote the subset $\{t \in [0, 2\pi) : e^{it} \in [e^{i\xi}, e^{i\xi'}]\}$; the intervals $[\xi, \xi')$, $(\xi, \xi']$ and (ξ, ξ') will also be defined similarly.

In Example 2.9, a direct computation shows that for $1 \le r, k \le n$,

$$\xi_{r+k} - \xi_r = \begin{cases} 2k\pi/n & \text{if } r + k \le n, \\ 2k\pi/n - 2\pi & \text{if } r + k > n. \end{cases}$$

Therefore, the set $\{\xi_1, \dots, \xi_n\}$ is k-regular and $\Lambda_k(A)$ is nonempty for $1 \le k < n/2$. Otherwise, the set $\{\xi_1, \dots, \xi_n\}$ is not k-regular and $\Lambda_k(A)$ is either empty or a singleton.

In the following, we need an alternate formulation of (1.1). For any $d \in \mathbb{R}$ and $\xi \in [0, 2\pi)$, consider the closed half plane

$$\mathcal{H}(d, \xi) = \{\mu \in \mathbb{C} : \mathrm{Re}\,(e^{-i\xi}\mu) \le d\},$$

and its boundary, which is the straight line

$$\mathcal{L}(d, \xi) = \partial\mathcal{H}(d, \xi) = \{\mu \in \mathbb{C} : \mathrm{Re}\,(e^{-i\xi}\mu) = d\}.$$

For $A \in M_n$, let $\mathrm{Re}A = (A + A^*)/2$. Then (1.1) is equivalent to

$$\Lambda_k(A) = \bigcap_{\xi \in [0, 2\pi)} \mathcal{H}(\lambda_k(\mathrm{Re}\,(e^{-i\xi}A)), \xi).$$

Theorem 3.3. *Suppose $\mathcal{P}$ is a non-degenerate polygon on $\mathbb{C}$. Then there is a normal matrix $A \in M_n$ such that $\Lambda_k(A) = \mathcal{P}$ if and only if there are $d_1, \dots, d_n \in \mathbb{R}$ and a k-regular set $\{\xi_1, \dots, \xi_n\}$ such that*

$$\mathcal{P} = \bigcap_{j=1}^{n} \mathcal{H}(d_j, \xi_j).$$

Notice that a necessary condition for the set $\bigcap_{j=1}^{p} \mathcal{H}(d_j, \xi_j)$ to be a non-degenerate polygon is that

$$\{e^{i\xi_1}, \dots, e^{i\xi_p}\} \text{ is 1-regular.} \tag{3.2}$$

To prove Theorem 3.3, we need some lemmas.

Lemma 3.4. *Given $A = \operatorname{diag}(a_1, \dots, a_n)$ and $1 \le m < n$. Suppose the eigenvalues $a_{m+1}, \dots, a_n$ are in $\Lambda_k(A)$ but not extreme points of $\Lambda_k(A)$. Then*

$$\Lambda_k(\operatorname{diag}(a_1, \dots, a_m)) = \Lambda_k(A).$$

Proof. It suffices to show that if a_n is in $\Lambda_k(A)$ but not an extreme point of $\Lambda_k(A)$, then $\Lambda_k(\operatorname{diag}(a_1, \dots, a_{n-1})) = \Lambda_k(A)$.

Suppose a_n satisfy the above assumption. Clearly, $\Lambda_k(\operatorname{diag}(a_1, \dots, a_{n-1}))$ is a subset of $\Lambda_k(A)$. On the other hand, for any $1 \le j_1 < \cdots < j_{n-k} \le n-1$, $\Lambda_k(A) \subseteq \operatorname{conv}\{a_{j_1}, \dots, a_{j_{n-k}}, a_n\}$. Since a_n is not an extreme point of $\Lambda_k(A)$, it follows that a_n lies in $\operatorname{conv}\{a_{j_1}, \dots, a_{j_{n-k}}, a_n\}$ but is not its extreme point. Therefore,

$$\operatorname{conv}\{a_{j_1}, \dots, a_{j_{n-k}}\} = \operatorname{conv}\{a_{j_1}, \dots, a_{j_{n-k}}, a_n\}.$$

Thus,

$$\begin{aligned}
\Lambda_k(A) &= \bigcap\{\operatorname{conv}\{a_{j_1}, \dots, a_{j_{n-k+1}}\} : 1 \le j_1 < \cdots < j_{n-k} < j_{n-k+1} \le n\} \\
&\subseteq \bigcap\{\operatorname{conv}\{a_{j_1}, \dots, a_{j_{n-k}}, a_n\} : 1 \le j_1 < \cdots < j_{n-k} \le n-1\} \\
&= \bigcap\{\operatorname{conv}\{a_{j_1}, \dots, a_{j_{n-k}}\} : 1 \le j_1 < \cdots < j_{n-k} \le n-1\} \\
&= \Lambda_k(\operatorname{diag}(a_1, \dots, a_{n-1})).
\end{aligned}$$

□

Lemma 3.5. *Suppose $\mathcal{P} = \cap_{j=1}^{m} \mathcal{H}(d_j, \xi_j)$ such that $0 \le \xi_1 \le \cdots \le \xi_m < 2\pi$ and for each $r = 1, \dots, m$, there are $1 \le j_1 < \cdots < j_k \le m$ such that $\xi_{j_1}, \dots, \xi_{j_k} \in (\xi_r, \xi_r + \pi)$. For every $n \ge m$, there are real numbers $\hat{d}_1, \dots, \hat{d}_n$ and a k-regular set $\{\hat{\xi}_1, \dots, \hat{\xi}_n\}$ such that $\mathcal{P} = \cap_{j=1}^{n} \mathcal{H}(\hat{d}_j, \hat{\xi}_j)$ and $\mathcal{P} \cap \mathcal{L}(\hat{d}_j, \hat{\xi}_j) \ne \emptyset$ for each $j = 1, \dots, n$.*

Proof. Set $\hat{\xi}_1 = \xi_1$, and for $s \in \{2, \dots, m\}$, let $\hat{\xi}_s = \xi_s$ if $\xi_{s-1} < \xi_s$. For the remaining values, we have $\xi_{s-1} = \xi_s$ and we can write that

$$\hat{\xi}_{s-t_1} = \xi_{s-t_1+1} = \cdots = \xi_s = \cdots = \xi_{s+t_2} < \hat{\xi}_{s+t_2+1}$$

for some $t_1 \geq 1$ and $t_2 \geq 0$. Let $\ell = \min\{j : \hat{\xi}_j > \hat{\xi}_{s-t_1} + \pi\}$, then we can replace ξ_{s+j} by $\hat{\xi}_{s+j} = \xi_{s+j} + \epsilon_j$ for sufficient small $\epsilon_j > 0$ for $j = -t_1 + 1, -t_1 + 2, \dots, 0, \dots, t_2$ such that

$$\hat{\xi}_{s-t_1} < \hat{\xi}_{s-t_1+1} < \cdots < \hat{\xi}_s < \cdots < \hat{\xi}_{s+t_2} < \min\{\hat{\xi}_\ell - \pi, \hat{\xi}_{s+t_2+1}\}.$$

After this modification, $\hat{\xi}_1, \dots, \hat{\xi}_m$ are distinct and $\{\hat{\xi}_1, \dots, \hat{\xi}_m\}$ is k-regular. If $n > m$, pick distinct $\hat{\xi}_{m+1}, \dots, \hat{\xi}_n \in [0, 2\pi) \backslash \{\hat{\xi}_1, \dots, \hat{\xi}_m\}$. Then $\{\hat{\xi}_1, \dots, \hat{\xi}_n\}$ also forms a k-regular set. Finally, let $\hat{d}_j = \max_{\mu \in \mathcal{P}} \operatorname{Re}\left(e^{-i\hat{\xi}_j}\mu\right)$ for $j = 1, \dots, n$. Clearly, we have $\mathcal{P} \cap \mathcal{L}(\hat{d}_j, \hat{\xi}_j) \neq \emptyset$ and $\mathcal{P} \subseteq \mathcal{H}(\hat{d}_j, \hat{\xi}_j)$ for all j. Since $\{\xi_1, \dots, \xi_m\}$ is a subset of $\{\hat{\xi}_1, \dots, \hat{\xi}_n\}$, hence $\mathcal{P} = \cap_{j=1}^m \mathcal{H}(d_j, \xi_j) = \cap_{j=1}^n \mathcal{H}(\hat{d}_j, \hat{\xi}_j)$. □

We can now present the proof of Theorem 3.3.

Proof of Theorem 3.3. Suppose $\mathcal{P} = \bigcap_{j=1}^n \mathcal{H}(d_j, \xi_j)$ is a non-degenerate polygon, where $\{\xi_1, \dots, \xi_n\}$ is k-regular. By Lemma 3.5, we may assume that $\mathcal{P} \cap \mathcal{L}(d_j, \xi_j) \neq \emptyset$ for all $j = 1, \dots, n$, and $0 \leq \xi_1 < \cdots < \xi_n < 2\pi$ such that condition (3.1) holds. For each $r = 1, \dots, n$, let

$$a_r = \frac{i}{\sin(\xi_{r+k} - \xi_r)} \left(e^{i\xi_r} d_{r+k} - e^{i\xi_{r+k}} d_r\right)$$

and $A = \operatorname{diag}(a_1, \dots, a_n)$. Then

$$\operatorname{Re}(e^{-i\xi_r} a_r) = d_r \quad \text{and} \quad \operatorname{Re}(e^{-i\xi_{r+k}} a_r) = d_{r+k}.$$

Note that $a_r \in \mathcal{L}(d_r, \xi_r) \cap \mathcal{L}(d_{r+k}, \xi_{r+k})$ is the vertex of the conical region $\mathcal{H}(d_r, \xi_r) \cap \mathcal{H}(d_{r+k}, \xi_{r+k})$, which contains $\mathcal{P}$. Therefore,

$$\operatorname{Re}(e^{-i\xi_r}(a_r - \mu)) \geq 0 \quad \text{and} \quad \operatorname{Re}(e^{-i\xi_{r+k}}(a_r - \mu)) \geq 0,$$

for all $\mu \in \mathcal{P}$. Since $\xi_{r+k} \in (\xi_r, \xi_r + \pi)$, we have

$$\operatorname{Re}(e^{-i\xi} a_r) \geq \max_{\mu \in \mathcal{P}} \operatorname{Re}(e^{-i\xi}\mu). \tag{3.3}$$

for all $\xi \in [\xi_r, \xi_{r+k}]$.

Let $\mu_j \in \mathcal{L}(d_j, \xi_j) \cap \mathcal{P}$ for $j = r, r+k$. As $\xi_{r+k} \in (\xi_r, \xi_r + \pi)$, we have $\mu_r = a_r - ie^{i\xi_r} b_r$ and $\mu_{r+k} = a_r + ie^{i\xi_{r+k}} c_r$ for some $b_r, c_r \geq 0$. Note that

$$\operatorname{Re}(e^{-i\xi}(\mu_r - a_r)) = b_r \sin(\xi_r - \xi) \geq 0, \quad \text{for all } \xi \in [\xi_r - \pi, \xi_r],$$

and

$$\operatorname{Re}(e^{-i\xi}(\mu_{r+k} - a_r)) = c_r \sin(\xi - \xi_{r+k}) \geq 0, \quad \text{for all } \xi \in [\xi_{r+k}, \xi_{r+k} + \pi].$$

Since $\{\xi_1, \dots, \xi_n\}$ is k-regular, it is easily seen that

$$[0, 2\pi) \setminus [\xi_r, \xi_{r+k}] = [\xi_r - \pi, \xi_r) \cup (\xi_{r+k}, \xi_{r+k} + \pi].$$

Therefore, for $\xi \in [0, 2\pi) \setminus [\xi_r, \xi_{r+k}]$, we have

$$\max\{\operatorname{Re}(e^{-i\xi}(\mu_r - a_r)), \operatorname{Re}(e^{-i\xi}(\mu_{r+k} - a_r))\} \geq 0.$$

Moreover, we have

$$\max\{\mathrm{Re}\,(e^{-i\xi}\mu_r), \mathrm{Re}\,(e^{-i\xi}\mu_{r+k})\} \geq \mathrm{Re}\,(e^{-i\xi}a_r). \tag{3.4}$$

Let $\xi \in [0, 2\pi)$. Then $\xi \in [\xi_s, \xi_{s+1})$ for some $s \in \{1, \ldots, n\}$. It follows that $\xi \in [\xi_r, \xi_{r+k}]$ for $r = s-k+1, \ldots, s$, and $\xi \in [0, 2\pi) \setminus [\xi_r, \xi_{r+k}]$ for other r. By (3.3) and (3.4),

$$\min_{r\in\{s-k+1,\ldots,s\}} \mathrm{Re}\,(e^{-i\xi}a_r) \geq \max_{\mu\in\mathcal{P}} \mathrm{Re}\,(e^{-i\xi}\mu) \geq \max_{r\notin\{s-k+1,\ldots,s\}} \mathrm{Re}\,(e^{-i\xi}a_r).$$

Thus, $\lambda_k(\mathrm{Re}\,(e^{-i\xi}A)) = \min_{r\in\{s-k+1,\ldots,s\}} \mathrm{Re}\,(e^{-i\xi}a_r)$ and so

$$\mathcal{P} \subseteq \mathcal{H}\left(\lambda_k(\mathrm{Re}\,(e^{-i\xi}A)), \xi\right).$$

Hence, $\mathcal{P} \subseteq \Lambda_k(A)$. Furthermore, if $\xi = \xi_s$, then $\mathrm{Re}\,(e^{-i\xi_s}a_s) = d_s$. Thus

$$\lambda_k(\mathrm{Re}\,(e^{-i\xi_s}A)) = \min_{r\in\{s-k+1,\ldots,s\}} \mathrm{Re}\,(e^{-i\xi_s}a_r) \leq d_s.$$

It follows that

$$\begin{aligned}\Lambda_k(A) &= \bigcap_{\xi\in[0,2\pi)} \mathcal{H}\left(\lambda_k(\mathrm{Re}\,(e^{-i\xi}A)), \xi\right) \\ &\subseteq \bigcap_{1\leq s\leq n} \mathcal{H}\left(\lambda_k(\mathrm{Re}\,(e^{-i\xi_s}A)), \xi_s\right) \subseteq \bigcap_{1\leq s\leq n} \mathcal{H}\left(d_s, \xi_s\right) = \mathcal{P}.\end{aligned}$$

Thus, $\mathcal{P} = \Lambda_k(A)$.

Next, we consider the converse. Suppose $A = \mathrm{diag}\,(a_1, \ldots, a_n)$ and $\Lambda_k(A) = \mathcal{P}$. By Lemma 3.4, we can remove the eigenvalues of A in $\Lambda_k(A)$ that are not extreme points of $\Lambda_k(A)$ to get $\tilde{A} \in M_m$ so that $\Lambda_k(A) = \Lambda_k(\tilde{A})$. If we can show that $\Lambda_k(\tilde{A}) = \cap_{j=1}^m \mathcal{H}(d_j, \xi_j)$ for a k-regular set $\{\xi_1, \ldots, \xi_m\}$, then Lemma 3.5 will ensure that $\Lambda_k(A) = \Lambda_k(\tilde{A}) = \cap_{j=1}^n \mathcal{H}(\tilde{d}_j, \tilde{\xi}_j)$ for some k-regular set $\{\tilde{\xi}_1, \ldots, \tilde{\xi}_n\}$.

For notational convenience, we assume that $A = \tilde{A}$ so that every eigenvalue is either an extreme point of $\Lambda_k(A)$ or not in $\Lambda_k(A)$.

Recall that in the proof of Theorem 2.3, $\mathcal{S}$ denotes the set of (r, s) such that $H(r, s)$ contains at least $n-k+1$ eigenvalues of A and

$$\mathcal{S}_r = \{s : (r, s) \in \mathcal{S}\}.$$

We first show that $\mathcal{S}_r \neq \emptyset$ for all r. As a_r is either an extreme point of $\Lambda_k(A)$ or not in $\Lambda_k(A)$, there is $\xi \in [0, 2\pi)$ such that $\mathrm{Re}\,(e^{-i\xi}a_r) \geq \lambda_k(\mathrm{Re}\,(e^{-i\xi}A))$. Thus, the closed half plane $\mathcal{H}(\mathrm{Re}\,(e^{-i\xi}a_r), \xi)$ contains at least $n-k+1$ eigenvalues of A. Let

$$\mathcal{W} = \mathrm{conv}\,\{a_t : a_t \in \mathcal{H}(\mathrm{Re}\,(e^{-i\xi}a_r), \xi)\}.$$

Then $\mathcal{W}$ is a convex set containing $\mathcal{P}$ and there is a vertex $a_s \in \mathcal{W}$ with $a_s \neq a_r$ such that $\mathcal{W} \subseteq H(r,s)$. Thus, $H(r,s)$ contains at least $n-k+1$ eigenvalues and hence $\mathcal{S}_r \neq \emptyset$.

Now by Theorem 2.3 and its proof,

$$\Lambda_k(A) = \bigcap_{\mathcal{S}_r \neq \emptyset} H(r,s_r) = \bigcap_{1 \le r \le n} H(r,s_r) = \bigcap_{1 \le r \le n} \mathcal{H}(d_r, \xi_r),$$

where $\xi_r = \arg(a_{s_r} - a_r) - \pi/2$ and $d_r = \mathrm{Re}\,(e^{-i\xi_r} a_r)$. We will prove the following.

Claim *For each $r = 1, \dots, n$, there are $1 \le j_1 < \cdots < j_k \le n$ such that $\xi_{j_1}, \dots, \xi_{j_k} \in (\xi_r, \xi_r + \pi)$.*

Once the claim is proved, Lemma 3.5 will ensure that $\Lambda_k(A) = \cap_{j=1}^n \mathcal{H}(\tilde{d}_j, \tilde{\xi}_j)$ for some k-regular set $\{\tilde{\xi}_1, \dots, \tilde{\xi}_n\}$.

To prove our claim, we first establish the following two assertions.

1. for any $\xi \in [0, 2\pi)$, if $\mathcal{H}(\mathrm{Re}\,(e^{-i\xi} a_r), \xi)$ contains at least $n-k+1$ eigenvalues of A, then either $\xi_r = \xi$ or $\sin(\xi_r - \xi) > 0$, and
2. the closed half plane $\{z \in \mathbb{C} : \mathrm{Re}\,(e^{-i\xi_r} z) \ge d_r\}$ contains at least k eigenvalues $a_{j_1}, \dots, a_{j_k}$ satisfying $\xi_{j_t} \neq \xi_r$.

To prove the two assertions, without loss of generality, we may assume that $a_r = 0$ and a_{s_r} lies in the positive imaginary axis. Then $d_r = 0$, $\xi_r = 0$ and

$$H(r, s_r) = H = \{z \in \mathbb{C} : \mathrm{Re}\,(z) \le 0\}.$$

Thus, the closed left half plane contains at least $n-k+1$ eigenvalues.

First we consider Assertion (1). Note that in terms of the set $\mathcal{S}_r$ associate with a_r in the proof of Theorem 2.3, Assertion (1) simply says that $\mathcal{S}_r$ cannot have any element in the interior of $H(r,s_r) = \mathcal{H}(d_r, \xi_r)$. Suppose $\mathcal{H}(\mathrm{Re}\,(e^{-i\xi} a_r), \xi)$ contains at least $n-k+1$ eigenvalues. By our assumption, $\mathcal{H}(\mathrm{Re}\,(e^{-i\xi} a_r), \xi) = e^{i\xi} H$.

Notice that the closed right half plane $-H$ can contain at most $n-k$ eigenvalues. Otherwise, $\Lambda_k(A)$ has to be a subset in $H \cap (-H) = i\mathbb{R}$, which contradicts the assumption that $\Lambda_k(A)$ is a non-degenerate polygon. From this argument, we first see that $\xi \neq \pi$.

Now suppose that $\xi \in (0, \pi)$. If the intersection of

$$H \cap (e^{i\xi} H) = \{z \in \mathbb{C} : \arg(z) \in [\xi + \pi/2, 3\pi/2]\}$$

does not contain any nonzero eigenvalue of A, then the closed half plane $-H$ will contain all the eigenvalues that are in $e^{i\xi} H$ and hence $-H$ has at least $n-k+1$ eigenvalues. But this is impossible by the previous argument. Thus, the intersection $H \cap (e^{i\xi} H)$ contains at least one nonzero eigenvalue of A. Take the nonzero eigenvalue a_s in $H \cap (e^{i\xi} H)$ such that a_s has the smallest argument $\arg(a_s) \in [\xi + \pi/2, 3\pi/2]$ among all the nonzero eigenvalues of A

in $H \cap (e^{i\xi}H)$. Let $\tau = \arg(a_s) - \pi/2$. Then $H(r,s) = e^{i\tau}H$ contains all the eigenvalues that are in $e^{i\xi}H$ and hence $H(r,s)$ contains at least $n-k+1$ eigenvalues. Then the property of a_{s_r} implies $a_{s_r} \in H(r,s)$ and so $H(r,s)$ contains the positive imaginary axis. As $\tau = \arg(a_s) - \pi/2 \in [\xi, \pi]$, this is possible only when $\tau = \pi$. But then $e^{i\tau}H = -H$, which contradicts our previous argument. Therefore, we must have $\xi \in (\pi, 2\pi]$ and hence Assertion (1) holds.

Next, turn to Assertion (2). Suppose that the closed half plane $-H$ contains eigenvalues $a_{j_1}, \dots, a_{j_{g+h}}$ with $\xi_{jt} = 0$ for $t = 1, \dots, g$, and $\xi_{jt} \neq 0$ for $t = g+1, \dots, g+h$. Then Assertion (2) holds if $h \geq k$. Fix a sufficiently small $\epsilon > 0$. We choose $1 \leq \ell \leq g$ so that

$$\mathrm{Re}\,(e^{-i\epsilon}a_{j_\ell}) = \max_{1 \leq t \leq g} \mathrm{Re}\,(e^{-i\epsilon}a_{j_t}).$$

Then $\{a_{j_1}, \dots, a_{j_g}\} \subseteq \mathcal{H}(\mathrm{Re}\,(e^{-i\epsilon}a_{j_\ell}), \epsilon)$. On the other hand, this closed half plane $\mathcal{H}(\mathrm{Re}\,(e^{-i\epsilon}a_{j_\ell}), \epsilon)$ also contains all eigenvalues of A that are in the left open half plane. Thus, this closed half plane $\mathcal{H}(\mathrm{Re}\,(e^{-i\epsilon}a_{j_\ell}), \epsilon)$ has at least $n-h$ eigenvalues of A. By Assertion (1), $\mathcal{H}(\mathrm{Re}\,(e^{-i\epsilon}a_{j_\ell}), \epsilon)$ has at most $n-k$ eigenvalues. Thus, we have $h \geq k$ and the assertion holds.

Now we prove our **claim**. By Assertion (2), for any $r = 1, \dots, n$, the closed half plane $\{z \in \mathbb{C} : \mathrm{Re}\,(e^{-i\xi_r}z) \geq d_r\}$ has at least k eigenvalues $a_{j_1}, \dots, a_{j_k}$ with $\xi_{jt} \neq \xi_r$. For each $t = 1, \dots, k$, let $h_t = \mathrm{Re}\,(e^{-i\xi_r}a_{j_t})$, then $h_t \geq d_r$ and $\mathcal{H}(d_r, \xi_r) \subseteq \mathcal{H}(h_t, \xi_r)$. Thus, the closed half plane $\mathcal{H}(h_t, \xi_r)$ contains at least $n-k+1$ eigenvalues. Now by Assertion (1), we see that

$$\sin(\xi_{j_t} - \xi_r) > 0 \quad \text{for } t = 1, \dots, k. \tag{3.5}$$

Thus, our **claim** is proved, and the result follows. □

By Theorem 3.3, we can deduce the following.

Theorem 3.6. *Let $\mathcal{P} = \bigcap_{j=1}^{p} \mathcal{H}(d_j, \xi_j)$ be a non-degenerate p-sided polygon, where $d_1, \dots, d_p \in \mathbb{R}$ and $\xi_1, \dots, \xi_p \in [0, 2\pi)$. The following two statements are equivalent.*

(I) *There is a $(p+q) \times (p+q)$ normal matrix A such that $\Lambda_k(A) = \mathcal{P}$.*
(II) *There are $\xi_{p+1}, \dots, \xi_{p+q} \in [0, 2\pi)$ such that $\{\xi_1, \dots, \xi_{p+q}\}$ is k-regular.*

Proof. Let $\mathcal{P} = \bigcap_{j=1}^{p} \mathcal{H}(d_j, \xi_j)$ be a non-degenerate p-sided polygon, where $d_1, \dots, d_p \in \mathbb{R}$ and $\xi_1, \dots, \xi_p \in [0, 2\pi)$.

Suppose (I) holds. By Theorem 3.3, there are $h_1, \dots, h_{p+q} \in \mathbb{R}$ and $\zeta_1, \dots, \zeta_{p+q} \in [0, 2\pi)$ such that $\{\zeta_1, \dots, \zeta_{p+q}\}$ is k-regular and

$$\bigcap_{j=1}^{p} \mathcal{H}(d_j, \xi_j) = \mathcal{P} = \Lambda_k(A) = \bigcap_{j=1}^{p+q} \mathcal{H}(h_j, \zeta_j).$$

Then $\{\xi_1, \dots, \xi_p\} \subseteq \{\zeta_1, \dots, \zeta_{p+q}\}$. Now we can take $\xi_{p+1}, \dots, \xi_{p+q} \in [0, 2\pi)$ so that $\{\xi_1, \dots, \xi_{p+q}\} = \{\zeta_1, \dots, \zeta_{p+q}\}$. Therefore, (II) holds.

Suppose (II) holds, namely, there are $\xi_{p+1}, \dots, \xi_{p+q}$ such that $\{\xi_1, \dots, \xi_{p+q}\}$ is k-regular. For $j = p+1, \dots, p+q$, define

$$d_j = \max_{\mu \in \mathcal{P}} \operatorname{Re}(e^{-i\xi_j}\mu).$$

Then $\mathcal{P} \subseteq \mathcal{H}(d_j, \xi_j)$ and so

$$\mathcal{P} = \bigcap_{j=1}^{p} \mathcal{H}(d_j, \xi_j) = \bigcap_{j=1}^{p+q} \mathcal{H}(d_j, \xi_j).$$

Then the result follows from Theorem 3.3. □

By Theorem 3.6, Problem 3.1 is equivalent to the following combinatorial problem.

Problem 3.7. Suppose $\{\xi_1, \dots, \xi_p\}$ is 1-regular. For $k > 1$, determine the smallest q so that $\{\xi_1, \dots, \xi_{p+q}\}$ is k-regular for some $\xi_{p+1}, \dots, \xi_{p+q} \in [0, 2\pi)$.

4. SOLUTIONS FOR PROBLEMS 3.1 AND 3.7

In this section, we give the solutions for Problems 3.1 and 3.7. Given a set of 1-regular complex units $\Pi = \{\alpha_1, \dots, \alpha_p\} \subseteq \Omega$, Problem 3.7 is equivalent to the study of smallest q so that $\{\alpha_1, \dots, \alpha_{p+q}\}$ is k-regular for some $\alpha_{p+1}, \dots, \alpha_{p+q} \in \Omega$. As shown in Assertion 1 later in this section, the existence of $\alpha_i, \alpha_j \in \Pi$ with $\alpha_j = -\alpha_i$ has implication on the size of a k-regular set Π. So, to deal with our problems, we always partition a set of complex units $\Pi = \{\alpha_1, \dots, \alpha_p\} \subseteq \Omega$ into $\Pi = \Pi_1 \cup \Pi_2$ with

$$\Pi_1 = \{\alpha_j \in \Pi : -\alpha_j \notin \Pi\} \quad \text{and} \quad \Pi_2 = \{\alpha_j \in \Pi : -\alpha_j \in \Pi\}. \tag{4.1}$$

Here either Π_2 is empty or Π_2 contains an even number of elements. For any finite set $\mathcal{S}$, denote the number of elements of $\mathcal{S}$ by $n(\mathcal{S})$. Suppose $n(\Pi_1) = r$ and $n(\Pi_2) = 2s$. Then $0 \le s \le p/2$ and $p = r + 2s$. To see the impact of the size of Π_1 and Π_2 on the solution of our optimization problem, let us consider the following example.

Example 4.1. Suppose $S_1 = \{1, w, w^2, w^3\}$ with $w = e^{2i\pi/5}$. Then $\alpha \ne -\beta$ for any two elements $\alpha, \beta \in S_1$ and adding w^4 to S_1 results in a 2-regular set. Suppose $S_2 = \{1, -1, i, -i\}$. Then we need to add at least two points, say, $z, -z \in \Omega \setminus S_2$, to get a 2-regular set.

In general, we have the following result allowing us to determine the minimum number of elements to be added to a finite set $\Pi \subseteq \Omega$ to get a k-regular set. We use the convention that a non-empty subset set of Ω is 0-regular in Theorem 4.2.

Theorem 4.2. *Given* $\Pi = \Pi_1 \cup \Pi_2$ *as defined in (4.1) with* $n(\Pi_1) = r$ *and* $n(\Pi_2) = 2s$. *Suppose* Π *is* 1*-regular. Then there is a set* $\Pi_3 \subseteq \Omega$ *with* $n(\Pi_3) = q$ *such that* $\Pi_1 \cup \Pi_2 \cup \Pi_3$ *is* k*-regular if and only if one of the following holds.*

(a) $k \geq r + s$ *and*

$$q \geq \ell^* := \begin{cases} 2k+1-p & \text{if } s = 0, \\ 2k+2-p & \text{if } s \neq 0. \end{cases}$$

(b) $k < r + s$ *and*

$$q \geq t^*,$$

where t^* *is the minimum* t *such that* Π_1 *contains a subset* $\hat{\Pi}_1$ *with* $n(\hat{\Pi}_1) = r - t$ *and* $\hat{\Pi}_1 \cup \Pi_2$ *being* $(k-t)$*-regular.*

Proof. Let $P = \{z \in \mathbb{C} : \operatorname{Im}(z) > 0\}$ be the open upper half plane. We divide the proof into five assertions.

Assertion 1. *Suppose* $S = \{\alpha_1, \ldots, \alpha_n\}$ *is* k*-regular. Then* $n \geq 2k+1$. *Furthermore, if there are distinct* i *and* j *such that* $\alpha_j = -\alpha_i$, *then* $n \geq 2k+2$.

Proof. For any $r \in \{1, \ldots, n\}$, each of the intervals $(\alpha_r, -\alpha_r)$ and $(-\alpha_r, \alpha_r)$ contains k elements of S. Thus, $n \geq 2k+1$. For the last statement, if we take $r = j$, then there are $2k$ elements in the intervals. Together with α_i and α_j, we see that $n \geq 2k+2$. The proof of the assertion is complete.

Assertion 2. *Suppose* $k \geq r+s$. *Then there is a set* $\Pi_3 \subseteq \Omega$ *with* $n(\Pi_3) = \ell^*$ *such that* $\Pi_1 \cup \Pi_2 \cup \Pi_3$ *is* k*-regular.*

Proof. Suppose $k \geq r + s$. Let Π' be a set containing $(k - r - s + 1)$ pairs of opposite points of the form $\{\alpha, -\alpha\}$ such that $\Pi' \cap \Pi$ is empty. Take $\Pi_3 = (-\Pi_1) \cup \Pi'$. Then

$$n(\Pi_3) = n(-\Pi_1) + n(\Pi') = r + 2(k - r - s + 1) = 2k + 2 - p.$$

Furthermore, the set $\Pi_1 \cup \Pi_2 \cup \Pi_3$ contains $k+1$ pairs of opposite points of the form $\{\alpha, -\alpha\}$ and hence it is k-regular. Thus, the result follows if $s \neq 0$.

Now suppose $s = 0$, i.e., $\Pi_2 = \emptyset$. Without loss of generality, we may assume that $1 \in \Pi_1$ and $-1 \in \Pi_3$. We now modify Π_3. We first delete the point -1 in Π_3. Then for all other points $\alpha \in \Pi_3$, we replace α by $e^{i\xi}\alpha$ with sufficiently small positive $\xi > 0$ if α lies in P, and with sufficiently small negative $\xi < 0$ if α lies in $-P$. Then we see that for every $\alpha \in \Pi_1 \cup \Pi_3$, αP still contains exactly k elements. Thus, $\Pi_1 \cup \Pi_3$ is k-regular. Furthermore, the modified set Π_3 has one fewer point, i.e., $n(\Pi_3) = 2k + 1 - p$. The proof of the assertion is complete.

Combining Assertions 1 and 2, we complete the proof of the case (a).

Assertion 3. *Suppose* $k < r+s$ *and there is a subset* $\hat{\Pi}_1$ *of* Π_1 *with* $n(\Pi_1) = r-t$ *such that* $\hat{\Pi}_1 \cup \Pi_2$ *is* $(k-t)$*-regular. Then there is a set* $\Pi_3 \subseteq \Omega$ *with* $n(\Pi_3) = t$ *such that* $\Pi_1 \cup \Pi_2 \cup \Pi_3$ *forms a* k*-regular set.*

Proof. As $\hat{\Pi}_1 \cup \Pi_2$ is $(k-t)$-regular, for every $\alpha \in \Omega$, the open half plane αP contains at least $k-t$ elements of $\hat{\Pi}_1 \cup \Pi_2$.

Without loss of generality, we assume $\hat{\Pi}_1 = \{\alpha_{t+1}, \dots, \alpha_r\} \subseteq \Pi_1$. Take $\alpha_{p+j} = -\alpha_j$ for $j = 1, \dots, t$ and let $\Pi_3 = \{\alpha_{p+1}, \dots, \alpha_{p+t}\}$. Then $(\Pi_1 \setminus \hat{\Pi}_1) \cup \Pi_3$ contains t pairs of opposite points of the form $\{\alpha, -\alpha\}$. Now for every $\alpha \notin (\Pi_1 \setminus \hat{\Pi}_1) \cup \Pi_3$, the open half plane αP contains at least $k-t$ elements of $\hat{\Pi}_1 \cup \Pi_2$ and at least t elements of $(\Pi_1 \setminus \hat{\Pi}_1) \cup \Pi_3$. Hence, αP contains at least k elements of $\Pi_1 \cup \Pi_2 \cup \Pi_3$. On the other hand, when $\alpha \in (\Pi_1 \setminus \hat{\Pi}_1) \cup \Pi_3$, the open half plane αP contains at least $k-1$ elements of $\Pi_1 \cup \Pi_2 \cup \Pi_3$.

By Assertion 1, $\hat{\Pi}_1 \cup \Pi_2$ has at least $2(k-t)+1$ elements and hence $\Pi_1 \cup \Pi_2 \cup \Pi_3$ contains at least $2k+1$ elements. Then for each $j = 1, \dots, t$, at least one of the open half plane, $\alpha_j P$ or $\alpha_{p+j} P (= -\alpha_j P)$, contains at least k elements while the other contains at least $k-1$ elements. As α_{p+j} lies in the boundary of both $\alpha_j P$ and $\alpha_{p+j} P$, we can perturb α_{p+j} so that both the open half planes $\alpha_j P$ and $\alpha_{p+j} P$ contain at least k elements of $\Pi_1 \cup \Pi_2 \cup \Pi_3$. Thus the assertion follows.

Assertion 4. *Suppose there is a set* $\Pi_3 \subseteq \Omega$ *with* $n(\Pi_3) = q$ *such that* $\Pi_1 \cup \Pi_2 \cup \Pi_3$ *is* k*-regular. Then there exist a subset* $\hat{\Pi}_1$ *of* Π_1 *and a set* $\tilde{\Pi}_3 \subseteq \Omega$ *with* $n(\hat{\Pi}_1) = r-1$ *and* $n(\tilde{\Pi}_3) = q-1$ *such that* $\hat{\Pi}_1 \cup \Pi_2 \cup \tilde{\Pi}_3$ *is* $(k-1)$*-regular.*

Proof. We claim that there are $\beta \in \Pi_3$ and $\gamma \in \Pi_1 \cup \Pi_3$ such that the set $(\Pi_1 \cup \Pi_2 \cup \Pi_3) \setminus \{\beta, \gamma\}$ is $(k-1)$-regular.

Suppose our claim is proved. If $\gamma \in \Pi_1$, then the result follows with $\hat{\Pi}_1 = \Pi_1 \setminus \{\gamma\}$ and $\tilde{\Pi}_3 = \Pi_3 \setminus \{\beta\}$. If $\gamma \in \Pi_3$, then the result follows with $\hat{\Pi}_1 = \Pi_1 \setminus \{\alpha\}$ for any $\alpha \in \Pi_1$ and $\tilde{\Pi}_3 = (\Pi_3 \cup \{\alpha\}) \setminus \{\beta, \gamma\}$.

It remains to prove our claim. If there is $\beta \in \Pi_3$ such that $-\beta \in \Pi_1 \cup \Pi_3$, then it is easy to see that the set $(\Pi_1 \cup \Pi_2 \cup \Pi_3) \setminus \{\beta, -\beta\}$ is $(k-1)$-regular. Thus, the claim follows by taking $\gamma = -\beta$.

We can assume that for each $\beta \in \Pi_3$, $-\beta \notin \Pi_1 \cup \Pi_3$. Notice that we may assume that $\Pi_3 \cap \Pi = \emptyset$. So we have $-\beta \notin \Pi_2$. Fixed a point $\beta \in \Pi_3$. Without loss of generality, we may assume that $\beta = -1$.

Furthermore, we have the following extra assumption when $\Pi_2 \neq \emptyset$. By Assertion 1, $\Pi_1 \cup \Pi_2 \cup \Pi_3$ contains at least $2k+2$ elements. Thus, either the upper open half plane or the lower open half plane contains at least $k+1$ elements. By the fact that a set $\mathcal{S}$ is k-regular if and only if the set $\bar{\mathcal{S}} = \{\bar{z} : z \in \mathcal{S}\}$ is k-regular, replacing $\Pi_1 \cup \Pi_2 \cup \Pi_3$ with $\bar{\Pi}_1 \cup \bar{\Pi}_2 \cup \bar{\Pi}_3$ if necessary, we may further assume that the upper open half plane contains at least $k+1$ elements if Π_2 is non-empty.

Let γ be the point in $\Pi_1 \cup \Pi_3$ such that $\arg(\gamma) \leq \arg(\alpha)$ for all $\alpha \in \Pi_1 \cup \Pi_3$. We show that $(\Pi_1 \cup \Pi_2 \cup \Pi_3) \setminus \{\beta, \gamma\}$ is $(k-1)$-regular.

Take any $\alpha \in (\Pi_1 \cup \Pi_2 \cup \Pi_3) \setminus \{\beta, \gamma\}$. Suppose $\alpha \in \beta P \cup \gamma P$. Then the open half plane αP can contain at most one of points β and γ. As the open half plane αP contains at least k elements of $\Pi_1 \cup \Pi_2 \cup \Pi_3$, αP contains at least $k-1$ elements of $(\Pi_1 \cup \Pi_2 \cup \Pi_3) \setminus \{\beta, \gamma\}$.

Now let $\omega_1, \ldots, \omega_t$ be the points in $(\Pi_1 \cup \Pi_2 \cup \Pi_3) \setminus (\beta P \cup \gamma P)$. By the choice of γ, $\{\omega_1, \ldots, \omega_t\} \subseteq \Pi_2$. Furthermore, all of them lie in the upper open half plane P. Therefore, we may assume that

$$0 < \arg(\omega_1) < \cdots < \arg(\omega_t) < \arg(\gamma) < \pi.$$

Clearly, each open half plane $\omega_j P$ contains at least k elements of $\Pi_1 \cup \Pi_2 \cup \Pi_3$. Suppose $\omega_j P$ contains exactly k elements. Notice that $\omega_j P$ contains $-\omega_1, \ldots, -\omega_{j-1}$, and β, which do not lie in the upper open half plane P. Thus, the intersection $P \cap (\omega_j P)$ contains at most $k-j$ elements and hence the upper open half plane P contains at most k elements only. But this contradicts to our assumption that the upper open half plane contains at least $k+1$ elements when $\Pi_2 \neq \emptyset$. Thus, every open half plane $w_j P$ contains at least $k+1$ elements of $\Pi_1 \cup \Pi_2 \cup \Pi_3$ and hence every open half plane $w_j P$ contains at least $k-1$ elements of $(\Pi_1 \cup \Pi_2 \cup \Pi_3) \setminus \{\beta, \gamma\}$. Therefore, $(\Pi_1 \cup \Pi_2 \cup \Pi_3) \setminus \{\beta, \gamma\}$ is a $(k-1)$-regular set and the assertion follows.

Assertion 5. *Suppose there is a set* $\Pi_3 \subseteq \Omega$ *with* $n(\Pi_3) = q$ *such that* $\Pi_1 \cup \Pi_2 \cup \Pi_3$ *is* k*-regular. Let* $t = \min\{k, q, r\}$*. Then there are subsets* $\hat{\Pi}_1$ *of* Π_1 *and* $\tilde{\Pi}_3$ *of* Ω *with* $n(\hat{\Pi}_1) = r-t$ *and* $n(\tilde{\Pi}_3) = q-t$ *such that* $\hat{\Pi}_1 \cup \Pi_2 \cup \tilde{\Pi}_3$ *is* $(k-t)$*-regular. Furthermore, if* $k < r+s$*, the set* $\hat{\Pi}_1 \cup \Pi_2$ *is also* $(k-t)$*-regular.*

Proof. The first part is clear by Assertion 4. Assume $k < r+s$. The last assertion is also clear when $t = q$ as $\tilde{\Pi}_3 = \emptyset$ in this case.

Suppose $t = k$. As $k < r+s$, either $r > k$ or $s > 0$. In both cases, $\hat{\Pi}_1 \cup \Pi_2$ is 0-regular. Finally, suppose $t = r$. Then $s > 0$. Notice that $\hat{\Pi}_1 \cup \Pi_2$ contains s pairs of opposite points of the form $\{\alpha, -\alpha\}$. Thus, the set $\hat{\Pi}_1 \cup \Pi_2$ is $(s-1)$-regular and hence $(k-r)$-regular as $k-r \leq s-1$. Then the result follows.

Combining the five assertions, we get the conclusion in Theorem 4.2. □

Remark 4.3. *Note that if condition (a) of the theorem holds, then the minimum number* ℓ^* *can be computed immediately. If condition (b) holds, one can check each* $r-t$ *element subset* $\hat{\Pi}_1$ *of* Π_1 *to see whether* $\hat{\Pi}_1 \cup \Pi_2$ *is* $k-t$ *regular, for* $t = 0, 1, \ldots$*. Thus, in a finite number of steps, one can determine the minimum number* t^**.*

The following proposition gives more information about the value ℓ^* and t^* in Theorem 4.2.

Proposition 4.4. *Using the notation in Theorem 4.2. If $k < r + s$, then the value t^* exists and satisfies*

$$\ell^* \le t^* \le \begin{cases} k & \text{if } (r,s) = (k+1,0) \text{ or } (k,1), \\ k-1 & \text{otherwise.} \end{cases}$$

Furthermore, $t^* = \ell^*$ if $p \le k+2$ with $s = 0$ or $p \le k+3$ with $s \ne 0$.

Proof. We first consider the case when $(r,s) = (k+1,0)$ or $(k,1)$. Take $t = k$ and any arbitrary subset $\hat{\Pi}_1$ of Π_1 with $n(\Pi_1) = r - t$. In both cases, $\hat{\Pi}_1 \cup \Pi_2$ is non-empty and hence is 0-regular. Thus, $t^* \le t = k$.

Now we assume that $(r,s) \notin \{(k+1,0),(k,1)\}$. Consider the case when $s \ge 2$. Take $t = \min\{k-1, r\}$ and an arbitrary subset $\hat{\Pi}_1$ of Π_1 with $n(\hat{\Pi}_1) = r - t$. Clearly, $\hat{\Pi}_1 \cup \Pi_2$ is $(s-1)$-regular and hence $(k-t)$-regular as $k - t = \max\{1, k-r\} \le s - 1$. Thus, we have $t^* \le t \le k-1$.

Next we consider the case when $s = 1$ and $r \ge k+1$. Let $\Pi_2 = \{\alpha, \ -\alpha\}$. Since Π is 1-regular, there exists $\alpha_1 \in (\alpha, \ -\alpha) \cap \Pi_1$ and $\alpha_2 \in (-\alpha, \ \alpha) \cap \Pi_1$. Take $t = k-1$. Choose any subset $\hat{\Pi}_1$ of Π_1 with $n(\hat{\Pi}_1) = r - t \ge 2$ such that $\alpha_1, \ \alpha_2 \in \hat{\Pi}_1$. Then $\hat{\Pi}_1 \cup \Pi_2$ is 1-regular. Then the result follows.

Finally consider the case when $s = 0$ and $r \ge k+2$. We may assume that $\Pi = \{e^{i\xi_j} : 1 \le j \le p\}$ with $0 = \xi_1 < \cdots < \xi_p < 2\pi$. Since Π is 1-regular, we can choose ℓ such that $\xi_\ell = \max\{\xi_j : 0 < \xi_j < \pi\}$. Then $S = \{\xi_1, \xi_\ell, \xi_{\ell+1}\}$ is 1-regular. Take $t = k-1$. Let $S \subseteq \hat{\Pi}_1 \subseteq \Pi_1$ with $n(\hat{\Pi}_1) = r - t \ge 3$. Then $\hat{\Pi}_1 \cup \Pi_2$ is 1-regular. □

By Theorem 4.2 and Proposition 4.4, we can answer Problems 3.1 and 3.7 and obtain some additional information on the solutions.

Theorem 4.5. *Let $\Pi = \{e^{i\xi_j} : 1 \le j \le p\}$. The optimal solution (minimum value) $\tilde{q}$ in Problem 3.7 is equal to ℓ^* or t^*, depending on case (a) or (b) of Theorem 4.2. Furthermore,*

$$\tilde{q} \le \max\{k-1, 2k+2-p\}. \tag{4.2}$$

For Problem 3.1, if a p-sided polygon $\mathcal{P}$ is expressed as $\mathcal{P} = \cap_{j=1}^{p} \mathcal{H}(d_j, \xi_j)$ for some $d_1, \dots, d_p > 0$, then the minimum dimension $\tilde{n}$ for the existence of a normal matrix $A \in M_{\tilde{n}}$ such that $\Lambda_k(A) = \mathcal{P}$ is equal to $p + \ell^*$ or $p + t^*$ depending on case (a) or (b) of Theorem 4.2. Furthermore,

$$\tilde{n} \le \max\{p+k-1, 2k+2\}. \tag{4.3}$$

Moreover, the inequalities in (4.2) and (4.3) become equalities if we let $\Pi = \{\alpha_1, \dots, \alpha_p\}$, where $(\alpha_1, \alpha_2, \alpha_3) = (1, i, -1)$ and $\alpha_4, \dots, \alpha_p$ lie in the open lower half plane.

Proof. The assertions on $\tilde{q}$ and $\tilde{n}$ are clear. To see the last assertion, we see that in order to get a k-regular set by adding q points to Π, we need to add at least $k-1$ points $e^{i\xi}$, with $0 < \xi < \pi$. If $2k+2-p > k-1$, then $p-3 < k$ and we need to add an extra $k-(p-3)$ points $e^{i\xi}$, with $\pi < \xi < 2\pi$, giving a total of $k-1+k-(p-3) = 2k+2-p$ points. This proves the equality in (4.2), from which the equality in (4.3) follows. $\square$

To close our paper, let us illustrate our results by the following example.

Example 4.6. Let the polygon $\mathcal{P} = \operatorname{conv}\{1, w, w^2, w^3, w^4, w^5, w^6, w^9\}$ with $w = e^{2\pi i/12}$, see the following.



The polygon $\mathcal{P}$

Then $\mathcal{P} = \bigcap_{j=1}^{8} \mathcal{H}(d_j, \xi_j)$ with $d_1 = \cdots = d_6 = \cos\frac{\pi}{12}$, $d_7 = d_8 = \cos\frac{\pi}{4}$, and

$$(\xi_1, \ldots, \xi_8) = \left(\frac{\pi}{12}, \frac{3\pi}{12}, \frac{5\pi}{12}, \frac{7\pi}{12}, \frac{9\pi}{12}, \frac{11\pi}{12}, \frac{15\pi}{12}, \frac{21\pi}{12}\right).$$

Thus,

$$\Pi = \{\alpha_1, \ldots, \alpha_8\} = \left\{e^{\frac{\pi i}{12}}, e^{\frac{3\pi i}{12}}, e^{\frac{5\pi i}{12}}, e^{\frac{7\pi i}{12}}, e^{\frac{9\pi i}{12}}, e^{\frac{11\pi i}{12}}, e^{\frac{15\pi i}{12}}, e^{\frac{21\pi i}{12}}\right\}.$$

In particular,

$$\Pi_1 = \left\{e^{\frac{\pi i}{12}}, e^{\frac{5\pi i}{12}}, e^{\frac{7\pi i}{12}}, e^{\frac{11\pi i}{12}}\right\} \quad \text{and} \quad \Pi_2 = \left\{e^{\frac{3\pi i}{12}}, e^{\frac{9\pi i}{12}}, e^{\frac{15\pi i}{12}}, e^{\frac{21\pi i}{12}}\right\},$$

i.e., $r = 4$ and $s = 2$. By Theorem 4.5 and Proposition 4.4, for $k \geq 5$, a $(2k+2) \times (2k+2)$ normal matrix A can be constructed so that $\Lambda_k(A) = \mathcal{P}$.

It remains to consider the cases for $k \leq 4$. Clearly, Π is 2-regular. Thus, a 8×8 normal matrix A_2 can be constructed so that $\Lambda_2(A_2) = \mathcal{P}$. However, Π is not k-regular for $k \geq 3$.

Now we consider the case $k = 3$. Then $\hat{\Pi}_1 \cup \Pi_2$ is 2-regular if $\hat{\Pi}_1 = \Pi_1 \setminus \{e^{\frac{5\pi i}{12}}\}$. Theorem 4.5 shows that there is a 9×9 normal matrix A_3 such that $\Lambda_3(A_3) = \mathcal{P}$. Indeed, following the proof of Assertion 3, we see that if $\Pi_3 = \{e^{\frac{18\pi i}{12}}\}$, $\Pi_1 \cup \Pi_2 \cup \Pi_3$ is 3-regular.

Finally, we turn to the case when $k = 4$. Notice that $\hat{\Pi}_1 \cup \Pi_2$ is 2-regular if $\hat{\Pi}_1 = \Pi_1 \setminus \{e^{\frac{5\pi i}{12}}, e^{\frac{7\pi i}{12}}\}$. Thus, Theorem 4.5 shows that there is a 10×10 normal matrix A_4 such that $\Lambda_4(A_4) = \mathcal{P}$.

In the following, we display the higher rank numerical ranges of A_2, A_3, and A_4. In the figures, the points "o" correspond to the vertices of the polygon while the points "$*$" correspond to the eigenvalues of the normal matrices.



$\Lambda_2(A_2) = \mathcal{P}$



$\Lambda_3(A_3) = \mathcal{P}$



$\Lambda_4(A_4) = \mathcal{P}$

Acknowledgement

This research began at the 2008 IMA PI Summer Program for Graduate Students, where the second author is a lecturer and the third author is a co-organizer. The support of IMA and NSF for the program is graciously acknowledged. The hospitality of the colleagues at Iowa State University is deeply appreciated.

References

[1] M.D. Choi, M. Giesinger, J. A. Holbrook, and D.W. Kribs, Geometry of higher-rank numerical ranges, Linear and Multilinear Algebra 56 (2008), 53-64.

[2] M.D. Choi, J.A. Holbrook, D. W. Kribs, and K. Życzkowski, Higher-rank numerical ranges of unitary and normal matrices, Operators and Matrices 1 (2007), 409-426.

[3] M.D. Choi, D. W. Kribs, and K. Życzkowski, Higher-rank numerical ranges and compression problems, Linear Algebra Appl. 418 (2006), 828-839.

[4] M.D. Choi, D. W. Kribs, and K. Życzkowski, Quantum error correcting codes from the compression formalism, Rep. Math. Phys., 58 (2006), 77–91.

[5] H.L. Gau, C.K. Li, and P.Y. Wu, Higher-Rank Numerical Ranges and Dilations, J. Operator Theory, to appear.

[6] R.A. Horn and C.R. Johnson, Topics in Matrix Analysis, Cambridge University Press, Cambridge, 1991.

[7] E. Knill and R. Laflamme, Theory of quantum error-correcting codes, Phys. Rev. A 55 (1997), 900-911.

[8] E. Knill, R. Laflamme, and L. Viola, Theory of quantum error correction for general noise, Phys. Rev. Lett. 84 (2000), 2525.

[9] D.W. Kribs, R. Laflamme, D. Poulin, and M. Lesosky, Operator quantum error correction, e-preprint. arXiv:quant-ph/0504189v3.

[10] S.R. Lay, Convex Sets and Their Applications, Pure and Applied Mathematics, John Wiley & Sons, Inc., New York, 1982.

[11] C.K. Li and Y.T. Poon, Quantum error correction and generalized numerical ranges, submitted. e-preprint http://arxiv.org/abs/0812.4772.

[12] C.K. Li, Y.T. Poon, and N.S. Sze, Higher rank numerical ranges and low rank perturbation of quantum channels, J. Math. Anal. Appl. 348 (2008), 843-855.

[13] C.K. Li, Y.T. Poon, and N.S. Sze, Condition for the higher rank numerical range to be non-empty, Linear and Multilinear Algebra, to appear. e-preprint http://arxiv.org/abs/0706.1540.

[14] C.K. Li and N.S. Sze, Canonical forms, higher rank numerical ranges, totally isotropic subspaces, and matrix equations, Proc. Amer. Math. Soc., 136 (2008), 3013-3023.

[15] M.A. Nielsen and I.L. Chuang, Quantum computation and quantum information, Cambridge, New York, 2000.

[16] H. Woerdeman, The higher rank numerical range is convex, Linear and Multilinear Algebra 56 (2008), 65-67.

Department of Mathematics, National Central University, Chung-Li 320, Taiwan
E-mail address: `hlgau@math.ncu.edu.tw`

Department of Mathematics, College of William & Mary, Williamsburg, VA 23185
E-mail address: `ckli@math.wm.edu`

Department of Mathematics, Iowa State University, Ames, IA 50051
E-mail address: `ytpoon@iastate.edu`

Department of Mathematics, University of Connecticut, Storrs, CT 06269
E-mail address: `sze@math.uconn.edu`